


How to Care for a Baby with Signs of Withdrawal



Use the Right Words

I was exposed to substances in utero. I am not an addict. And my parent may or may not have a Substance Use Disorder (SUD).




Treat Us as a Dyad

Parents and babies need each other. Help us bond. Whenever possible, provide my care alongside my parents and teach them how to meet my needs.




Support Rooming-In

Babies like me do best in a calm, quiet, dimly-lit room where we can be close to our caregivers.



Promote Kangaroo Care

Skin-to-skin care helps me stabilize and self-regulate. It helps relieve the autonomic symptoms associated with withdrawal, promotes bonding, and helps me sleep.



Try Non-Pharmacological Care

Help me self-soothe. Swaddle me snugly in a flexed position that reminds me of the womb. Offer me a pacifier to suck on. Protect my sleep by "clustering" my care.



Provide Lactation Support

Human milk is important to my gastrointestinal health and breastfeeding is recommended when my parent is HIV-negative and receiving medically-supervised care. Help my family reach our pumping and feeding goals.



Treat My Symptoms

If I am experiencing signs of withdrawal that make it hard for me to eat, sleep, and be soothed, create a care plan to help me wean comfortably.



Academy of Perinatal
Harm Reduction

www.perinatalharmreduction.org



www.nationalperinatal.org